

Alaska Earthcover Mapping



History of Earth Cover Mapping in Alaska

- late 1980's with Ducks Unlimited, Canada
- 1992 begin long-range outlook
- 1997 five-year plan
- 2005 over 151 million acres completed and more in progress.
- Earth Cover Classification System has approval of <u>Alaska Geographic Data Committee</u> and is being used in Canada, circumpolar arctic and Ukraine, accepted by 26 agencies, NGOs and conservation groups.

Basic Layers Within a Functional GIS



Earthcover

Hydrography

Digital Elevation Model

Transportation

Land Status

Why Alaska is Different!

- Large remote land base
- Limited dollars to gather traditional information on vegetation, soils, water, etc.
- Needed cost effective methodology

- Landscape level information for wildlife, subsistence, recreation, development, etc.
- Integration with other digital data
- Landscape level fire management

What We do!

- Ducks Unlimited and BLM developed a standardized approach to earth cover that was adopted by Alaska Geographic Data Committee and other agencies.
- Intensive field effort to collect training and accuracy assessment sites, essential for final map products.
- Continuing refinement of the technology for processing Landsat TM (30 meter).
- Each acre is now about \$ 0.05-.08!!!

ALASKA EARTH COVER CLASSIFICATION

FIELD WORK AND PROCESSING







PRE-FIELD WORK


- Image Pre-processing, selection of imagery, selection of training sites.
- Planning and Field Logistics



Image Pre-Processing

- Image selection/ordering
- QA/QC imagery
- Remove clouds from images
- Select field sites
- Produce plots of images with field sites


QC Imagery and Remove Clouds




Clouds/Haze

Shadows

Bad Pixels




Satellite Mapping







- Light reflected by vegetation
- Statistically isolate groups or "classes"

Select field Asites super vise of lales street in the range of land the single street in the single single



Select Field Sites





Plot Field Maps and Field Forms

Peu 43097					Gul kana	Fie	d For	m				
Yr Prob	cl Ce	- - 36)							Z3+ bs.Level Obs.Time:		г н	
Digital Phot	Sanda F	Photo		LAT (OPS)	Deck-	ulber	***		rowers	PS)	DechulDegrees	
%Slope (A	Jg)	Eleu	Aspect	N	NE E	7	SE	S	SW	JOJ.	NW	Flat
Aueroge Dis		32250	000000	1 1224	360 550	0.05	(22	5894	62.054.7	(Or	arer Weesland has d	HARONA
Be Ween St	ms:	10:19	15-27	20-3	i Chal-Lichens	-30		0-36	35 -40	-		
Fore	at I	Fore ct	- 0	Bhrub	Hertzgegu	_	Hertson	poor	Aquain Ve	own for	Barren	Othe
Closed Needless		Open Decido		Tal	Uchen		Dry Sedge		Aguaic Bed		Sparce Vég	Oh
Open Needlebear		Closed Mile		AZAL LOW	Moss		Dry Gramfrold		Bregent		Rock/Gesuel	=,0,0%
Woodland Needleben'		Open Mixed To		s sock Low Well Sedge / C		Gr	Dry Sedge/Gr		Snow/tz		IA.d/SIVSard	
Widdle Middle Buchen				Low-Lictery (Vile) Forts		ij. T			Turbidly	Valer		
Closed Dec	Mour		De	erflichen	Tussode-lid	en			Clear 9	lbbr .		
96CBV	Height TREES						њсо		нерез	RUCBO	nnet	
		NUM E Spr		Picesglauca				Parts G		19990 000200.		
3	Vi.	Black Sp.	-	Picsamatana				Savinge		Saxitage spp.		\neg
		Aspen		Populus fremulaities				Veth		As largedus um bellatus		
3	57	Blich		Be Uspanyri era				Horse talls		Enulse Limison		\neg
		Balsam Poplar		Populus bekennites				Fireweed		Epilodium angus Italium		
		Larch, Tanerack		Latixlatida				1		1		-1
	7			· C				1		18		=
96000	Height	8	H FRU EXB	21								\neg
		UNBOW		Salixapp.				Bryot	ď			
8		Alder		Ainus calspa				Moss				\neg
8		DwarfArc	ic Birch	Be Litanana	glandulosa			Ucher	n			
		Blueberry		Vacdrium uilginosum		\Box						\neg
		Low Bush Cranterry		Vacdrium disidea		9	96Cov AC		AQU	ATIC		
6		Crowteny		Empe kum rilgrum			Water Uly		Nupter polysepatum			
			Cirqueidi		Potrataspp.			Pordverd		Polimege brispp.		
E 1		Bearteny		Arios imphylos uumurst				Bultroup		Ramunodus spp.		
6 1		Mounish Auens		Dryas spp.				Mare's Tal		Hippuris spp.		
		Mountain Bell Heather		Cassique le lagore				Bucktean		Menyanites Mótala		
£ 1		Labeator Tea		Ledum pakis le				Warsh Ruefrger		Potraliapalus its		
8		Rose	9-971	Rosaladoula	mis .			Water	Sedge	Carexa	draft	_
		Leaterear		Cham.calyouals		-						-
				ŝ						1		
12.45(15)						9	6CBV			GETATE	D)	
96Cov	4000	HEREACE	OUB		-	-			Water			_
G.		«Graminoid	200			-			d Unbaler			_
	Pos	T				_		Shows		(drde	-	
-		ini Orass	Calamagros is canadarsi						3)VSard	(drde		_
-	-			tophorum spp.		-		Grate	l/Rock	(circle	are)	_
	-	Bedge Carexspp,				_		+		+-		
-				ke apre		_		100	KEKES.	1		
-	Benig	lenigass Agros is bon		ezas				-	Subloid % Couer			
_	_				_			GRA	D TOTAL 9	COVER	W.	
			1		1 2	_						



Customized data collection form

Pre-selected field sites

Field Work Planning

- Designate field crews
- Procurement of Aircraft
- Determine field camp locations
- Determine optimal sites for fuel locations
- Determine amount of fuel needed

Field Planning Map

TIEKEL PROJECT PLANNING MAP 25 Mile Fuel Buffer 50 Mile Fuel Buffer

FIELD CREW

Navigator:

Recorder:

Vegetation Caller:

Alternate:

Lori Scott

Dan

Mike

<u>Fuel</u>

Tiekel 500 gal. Glennallen 1000 gal.

FIELD WORK





Crew Roles/Responsibilities

Vegetation Caller




Navigator/Image Processor




Crew Roles/Responsibilities


Recorder/Alternate









Field Form input into DUFF Database





DUFF Data Entry Application




POST-FIELD WORK

- Field Site Database Quality Check
- Digitize New Field Sites
- Separate Accuracy Assessment Sites
- Classification
- Mosaic With Adjacent Classifications
- Final Products

Supervised Classification



TIEKEL PROJECT



Classified Map


Landsat TM Image

- **■** Closed Needleleaf
- Open Needleleaf
- Woodland Needleleaf
- Closed Deciduous
- Open Deciduous
- Closed Mixed Ndl./Dec.
- **■** Open Mixed Ndl./Dec.

- **Tall shrub**
- Low Shrub
- **Dwarf Shrub**
- Wet Graminoid
- Aquatic Bed
- **Emergent Vegetation**
- Clear Water

- Turbid Water.
- Ice/Snow
- **■** Rock/Gravel
- Sparse Vegetation
- Agriculture
- **■** Terrain Shadow

Mosaic Classifications




Existing Classification

+

New Classification


Mosaic












ALASKA EARTH COVER INITIATIVE COUCKS UNLIMITED








ALASKA EARTH COVER INITIATIVE COUCKS UNLIMITED











Canada






ALASKA EARTH COVER INITIATIVE COUCKS UNLIMITED






2006 PROJECT STATUS




BAER Earth Cover Project






ALASKA EARTH COVER INITIATIVE COUCKS UNLIMITED









Satellite imagery used for the BAER mapping project

BAER Project Area




BAER Project Mosaic



Uses for Management

- Wildlife and Waterfowl Habitat Selection
- Habitat Modeling
- Forest Fire History / Fuels Modeling
- Change Detection
- Environmental Impact Statements/CCP
- Road/Pipeline/Utility Placement
- Successional Vegetation Changes
- LandFire
- FRCC

Minto Flats - Golden Valley Electric Association Landcover Classification






Fire Fuels Modeling



Direct crosswalk from Earth Cover classifications to the Canadian **Forest Fire Behavior** Prediction Model




Kanuti NWR

Mologed Stuff Mymagg 99 July 2, 1999

- Fire Boundaries
- Refuge Boundary
 - River Survey
 - Transects

Digital Earth Cover of the NPR-A Arctic Coastal Plain





Focal Species

for resource selection analyses



Canada Goose



Brant



White-fronted Goose



Oldsquaw




Steller's Eider



King Eider

Waterfowl Densities





Earth Cover Selection Patterns

sedge/grass tussock tundra moss/lichen dwarf shrub low shrub tall shrub urbid water clear water dunes/sand sparse veg

Brant
Canada Goose
King Eider
Oldsquaw
Spectacled Eider
WF Goose
Steller's Eider

SELECTION KEY

strongly for

moderately for








Summary Values of Satellite Mapping



- Large-scale perspective and efficient value (\$.05-.08/acre)
- Earth cover data and protocol allows for repetition and change detection analyses over time
- Models can be refined and different scenarios tested, especially for monitoring and evaluation
- Field data collection contributes to Highly Accurate Final Products
- Continued work contributes to State Wide Earth Cover Data Set.

2006 PROJECT STATUS



Many thanks to our cooperators











